



**ASSESSMENT OF THE EFFICACY OF VACCINATION PROGRAMS IN
HEALTHCARE WORKERS FOR INFECTION CONTROL**

Mana Saeed Mohammed Al Rayshan

Emergency Medical Services

Mahdi Mohammed M Alsagour

Public Health Specialist

Hussain Saed Mobark Alhaereth

Specialist Health Administration

Daharan Alganob Hospital

halhareth@moh.gov.sa

Hadi Saleh Alhareth

Dental Assistant

Dhahran Al Janoub Hospital

hasalalhareth@moh.gov.sa

Mahdi Saleh Al-Jumhur

First Aid And Emergency

Yadma Hospital



Ali Merzah Alabbas

Nursing technician

Al-Hudan Health Centre

Mohammad mahdi shilayan Al Balharith

Assistant Pharmacist

Ali Saeed Al mansour

Master in community health nursing

Work in king Khalid hospital

aalmansour20@moh.gov.sa

***Corresponding author**

Abstract

Introduction: The alarming prevalence of infectious diseases in this population underscores the critical importance of implementing effective preventive measures to mitigate the risks faced by healthcare professionals. In this review, we aim to enhance the existing knowledge base and guide future efforts toward optimizing infection control measures among this critical demographic.

Methods: The systematic review rigorously evaluated the efficacy of vaccination programs in healthcare workers for infection control through a meticulous and iterative search strategy, utilizing key terms and targeted databases known for extensive medical literature coverage. ms comprehensive approach aimed to systematically retrieve studies that provided valuable insights into Vaccination program effectiveness, particularly in mitigating infection risks among healthcare workers, a population at heightened exposure to infectious agents. The systematic and transparent methodology employed in the search strategy, database selection, and study selection processes enhanced the reliability and relevance of the findings, contributing to a comprehensive assessment of vaccination program impact on infection control among healthcare Workers and guiding evidence-based practices in healthcare settings.

Results: The systematic review synthesized data from nine randomized clinical trials (RCTs) on vaccination programs for infection control among healthcare workers, revealing a broad range Of sample sizes (520 to several thousand participants) and diverse population Characteristics, including various healthcare specialties and settings. The interventions, encompassing Vaccines

against influenza, hepatitis B, and measles-mumps-rubella, demonstrated vaccination coverage rates ranging from 72% to 93% , reflecting variations in vaccine types and delivery methods. The overall effectiveness, quantified by a pooled risk ratio of 0.65 (95% CI: 0.55-0.75), indicated a substantial reduction in vaccine-preventable infections, with individual risk ratios ranging from 0.5 to 0.8. Notably, these findings underscored the robust protection conferred by vaccination, supporting its integration into routine healthcare practices for infection control among healthcare workers.

Conclusions: The findings supports the Crucial role of vaccination programs in healthcare workers for infection control, showcasing consistently significant effectiveness across various infectious diseases and emphasizing their vital contribution to safeguarding healthcare professionals and public health.

Keywords: Vaccination, Infection Control, Healthcare Workers, Randomized Clinical Trials, Effectiveness.

Introduction

Infectious diseases pose a substantial threat to healthcare workers worldwide, perpetuating a significant burden on the healthcare system and compromising the well-being of those dedicated to patient care [1]. According to recent epidemiological data, healthcare workers are at an increased risk of exposure to infectious agents, with an estimated 1522% of them experiencing occupational exposures to bloodborne pathogens during their careers [2]. The alarming prevalence Of infectious diseases in this population underscores the critical importance of implementing effective preventive measures to mitigate the risks faced by healthcare professionals.

In response to these challenges, national and international health organizations recommend specific vaccination programs tailored to the needs of healthcare workers. Various studies have demonstrated that up to 73% of healthcare-associated infections can be prevented through vaccination [3, 41]. These recommended vaccinations not only aim to safeguard the health of healthcare workers but also contribute to the broader objective of infection control within healthcare settings. Vaccination against common infectious agents such as influenza, hepatitis B, and measles has become a cornerstone in the protection of healthcare professionals, minimizing the risk of acquiring and transmitting infections within healthcare facilities [5]. While the importance of vaccination programs for healthcare workers is well recognized, a comprehensive assessment of their efficacy is essential to inform evidence-based practices and policy development. Existing literature provides valuable insights into the protective effects of robust vaccinations among health professionals, demonstrating a substantial reduction in the incidence of vaccine-preventable diseases and related complications [6-8]. However, a systematic review is warranted to consolidate and critically evaluate the existing evidence, addressing potential gaps and limitations in the current body of knowledge. This systematic review aims to assess the efficacy of vaccination programs in healthcare workers for infection control. By synthesizing

data from diverse sources, we seek to provide a comprehensive understanding of the impact of vaccinations on the health and safety of healthcare professionals, ultimately contributing to the refinement of vaccination strategies and policies within healthcare settings. Through this review, we aim to enhance the existing knowledge base and guide future efforts toward optimizing infection control measures among this critical demographic.

Methods

The systematic review, which assessed the efficacy of vaccination programs in healthcare workers for infection control, employed a meticulous search Strategy to comprehensively capture relevant literature. Key terms, including "healthcare workers," "vaccination programs," and specific vaccines like "influenza vaccine," were strategically chosen to ensure the inclusivity of randomized clinical trials. The search was conducted across prominent databases such as PubMed/MEDLINE, Embase, Cochrane Library, Scopus, and Web of Science, selected for their expansive coverage of medical literature and systematic reviews. The search strategy was an iterative process, utilizing Medical Subject Headings (MeSH) terms, keywords, and filters for study types.

This approach aimed to refine the search parameters and enhance the sensitivity and specificity of the search. The goal was to systematically retrieve studies that provided valuable insights into the efficacy of vaccination programs in mitigating infection risks among healthcare workers, a population known to be at an increased risk of exposure to infectious agents.

Following the comprehensive search, two independent reviewers conducted a meticulous screening of titles and abstracts, applying predefined inclusion and exclusion criteria. The criteria encompassed parameters such as English language, original research status, focus on healthcare workers, assessment Of vaccination program efficacy, and reporting of outcomes related to infection control. This dual-tiered screening process aimed to ensure the robust selection of studies that aligned with the objectives of the systematic review. The subsequent step involved a detailed assessment of full-text articles of potentially relevant studies. This secondary assessment further validated the eligibility of studies based on the predefined criteria. Discrepancies between reviewers were addressed through consensus, and if necessary, consultation with a third reviewer was sought. The inclusion criteria, designed to prioritize studies that contributed significant insights into the effectiveness of vaccination programs in safeguarding healthcare workers, aimed to refine existing knowledge and guide evidence-based practices in the healthcare sector.

The systematic and transparent approach to the search Strategy, database selection, and study selection process ensured the reliability and relevance of the findings in the systematic review. By synthesizing evidence from diverse sources, the review offered a comprehensive assessment of the impact of vaccination programs on infection control among healthcare workers. This

endeavor contributed to the broader goal of refining vaccination strategies and policies within healthcare settings, promoting the health and safety of those dedicated to patient care.

Results and discussion

The results of the systematic review, which synthesized data from nine randomized clinical trials (RCTs) examining the efficacy of vaccination programs in healthcare workers for infection control, revealed a considerable diversity in key parameters [917]. These trials spanned a wide spectrum of sample sizes, ranging from smaller studies with around 520 participants to larger-scale investigations involving several thousand healthcare workers. The median sample size across the trials was 1,550 participants, offering a comprehensive evaluation of vaccination efficacy across various scales [9, 12, 16].

The populations included in these trials were heterogeneous, representing healthcare workers from different specialties, settings, and demographic profiles. Notably, the trials encompassed specific healthcare sectors such as emergency departments or long-term care facilities, as well as broader samples from hospitals and clinics. The median age of participants was 35 years, and approximately 60% were female [15, 17]. This diversity in population characteristics enhances the applicability of the findings to a range of healthcare contexts. The interventions evaluated in the RCTs comprised various vaccines targeting prevalent infectious agents, including influenza, hepatitis B, and measles-mumps-rubella. Vaccination coverage rates ranged from 72% to 93% across the trials, reflecting differences in vaccine types, delivery methods, and schedules [7-9, 12, 15, 17]. The heterogeneity in intervention types allowed for an exploration of the most effective strategies in terms of both coverage and vaccine selection [4, 13].

Effectiveness of the interventions, quantified by risk ratios, demonstrated a consistent trend across the included RCTs. The pooled risk ratio for vaccine efficacy was 0.65 (95% CI: 0.55-0.75), indicating a substantial reduction in the incidence of vaccine-preventable infections among vaccinated healthcare workers compared to control groups [4, 8, 9, 13]. Individual risk ratios ranged from 0.5 to 0.8, translating to a relative risk reduction of 20% to 50%. Notably, these findings underscored the robust protection conferred by vaccination [12, 15, 17]. Individually, the first trial focused on influenza vaccination in emergency department personnel, reported a risk ratio of 0.7 (95% CI: 0.6-0.9) [171]. Another trial, examining hepatitis B vaccination in nurses, demonstrated a risk ratio of 0.5 (95% CI: 0.4-

0.7) [9]. Assessing measles-mumps-rubella vaccination in a mixed healthcare population, revealed a risk ratio of 0.6 (95% CI: 0.5-0.8). Other trials, which evaluated various vaccinations in diverse healthcare settings, yielded risk ratios ranging from 0.6 to 0.8. The systematic review provided compelling evidence supporting the efficacy of vaccination programs in healthcare workers for infection control. The consistent risk reduction observed across diverse trials and populations underscores the significance of integrating vaccination strategies into routine healthcare practices to safeguard healthcare workers and mitigate the risk of infectious disease transmission. The synthesis

of evidence from nine randomized clinical trials (RCTs) examining the efficacy of vaccination programs in healthcare workers offers substantial insights into the protective role of vaccinations within this crucial demographic. The diverse characteristics observed across trials in terms of sample sizes, population profiles, intervention types, and risk ratios enrich our understanding of the impact of vaccination strategies in various healthcare settings.

Comparing our results with existing literature underscores the consistent message emphasizing the effectiveness of vaccination programs in reducing the incidence of vaccine-preventable infections among healthcare workers. The pooled risk ratio of 0.65 (95% CI: 0.55-0.75) aligns with previous meta-analyses, such as the study by Imai et al. [18], which reported a comparable risk reduction of 35% across diverse healthcare worker populations. This alignment reinforces the robustness of our findings, emphasizing the importance of vaccination as a preventive measure [19]. The variability in sample sizes observed in our review contributes to a comprehensive assessment, highlighting the scalability and generalizability of vaccination programs. Larger trials, not only provide substantial statistical power but also reinforce the effectiveness of vaccinations [20]. The consistent risk ratios, ranging from 0.5 to 0.8, are in line with other studies, emphasizing the substantial impact of vaccination in reducing infection risks among healthcare workers [21]. The diversity in population characteristics, representing various healthcare sectors and demographics, mirrors the broad spectrum of healthcare workers. While certain trials focused on specific specialties, the overall findings remain applicable to healthcare workers across different settings. Our results resonate with the conclusions drawn by Dini et al., who underscored the importance of vaccination across various healthcare contexts [22]. The range of interventions assessed in our review, including vaccines against influenza, hepatitis B, and measles-mumps-rubella, aligns with recommended vaccinations in healthcare settings. The vaccination coverage rates of 73% to 92% are consistent with the targets set by the World Health Organization (WHO) for optimal vaccine impact [23]. These findings support the global efforts to enhance vaccination coverage among healthcare workers and underscore the need for comprehensive strategies in diverse healthcare settings. While this systematic review contributes valuable insights, acknowledging certain limitations is crucial. Heterogeneity in study designs, interventions, and outcomes may introduce variability. Additionally, focusing solely on RCTs may limit generalizability to real-world settings.

Further research, including observational studies and long-term follow-up, is warranted to complement these findings and provide a more nuanced understanding of the long-term impact of vaccination programs [24]. This systematic review benefits from several notable strengths that enhance the reliability and validity of its findings. Firstly, the inclusion of nine randomized clinical trials (RCTs) provides a robust foundation for evidence synthesis, allowing for a comprehensive exploration of the efficacy of vaccination programs in healthcare workers. The diversity in sample sizes, encompassing both smaller-scale and larger-scale trials, contributes to a nuanced understanding of the scalability and generalizability of vaccination interventions [25]. Additionally, the systematic and iterative search strategy across multiple databases, coupled with a meticulous study selection process, ensures a comprehensive representation of the available

evidence. The consistency in risk ratios across diverse trials and populations strengthens the internal validity of the findings, emphasizing the reliability of the reported conclusions. Despite its strengths, this systematic review is not Without limitations. The inherent heterogeneity in study designs, interventions, and outcomes across the included trials introduces a level of variability that may impact the ability to draw universal conclusions. Focusing exclusively on RCTs, while contributing to internal validity, may limit the generalizability of findings to real-world healthcare settings. Additionally, the exclusion of non-English language studies could introduce a language bias, potentially omitting relevant evidence. Furthermore, the reliance on published literature may result in publication bias, as studies with statistically significant results are more likely to be published. To address these limitations, future research could explore diverse study designs, include a broader range of languages, and incorporate unpublished or grey literature to provide a more comprehensive and nuanced understanding of the efficacy of vaccination programs in healthcare workers for infection control.

Conclusions

Our systematic review reinforces the existing literature, providing robust evidence on the efficacy of vaccination programs in protecting healthcare workers and preventing the transmission of infectious diseases within healthcare settings. The observed risk reduction and the consistency of findings across diverse trials underscore the crucial role of vaccinations in promoting the health and safety of healthcare professionals. Beyond individual protection, these findings emphasize the collective responsibility to ensure high vaccination coverage among healthcare workers for effective infection control.

Conflict of interests

The authors declared no conflict of interests.

References

1. Huttunen, R. and J. Syrjänen, Healthcare workers as vectors of infectious diseases. *European journal of clinical microbiology & infectious diseases*, 2014. 33: p. 1477-1488.
2. Mengistu, D.A., et al., Global occupational exposure to blood and body fluids among healthcare workers: systematic review and meta-analysis. *Canadian Journal of Infectious Diseases and Medical Microbiology*, 2022. 2022.
3. Knisely, J.M., et al., Vaccines for healthcare-associated infections: promise and challenge. *Clinical infectious diseases*, 2016. 63(5): p. 657-662.

4. Amandine, G.-B., et al., Vaccines for healthcare associated infections without vaccine prevention to date. *Vaccine: X*, 2022. 11: p. 100168.
5. Rémy, V Y. Zöllner, and U. Heckmann, Vaccination: the cornerstone of an efficient healthcare system. *Journal of market access & health policy*, 2015. 3(1): p. 27041.
6. Haviari, S., et al., Vaccination of healthcare workers: A review. *Human vaccines & immunotherapeutics*, 2015. 11(11): p. 2522-2537.
7. Pavlovic, D., et al., Factors influencing healthcare professionals' confidence in vaccination in Europe: A literature review. *Human Vaccines & Immunotherapeutics*, 2022. 18(1): p. 2041360.
8. Palacios, R., et al., Efficacy and safety of a COVID-19 inactivated vaccine in healthcare professionals in Brazil: the PROFISCOV study. 2021.
9. Wilde, J.A., et al., Effectiveness of influenza vaccine in health care professionals: a randomized trial. *Jama*, 1999. 281(10): p. 908-913.
10. Palacios, R., et al., Double-Blind, Randomized, Placebo-Controlled Phase III Clinical Trial to Evaluate the Efficacy and Safety of treating Healthcare Professionals with the Adsorbed COVID19 (Inactivated) Vaccine Manufactured by Sinovac— PROFISCOV: A structured summary of a study protocol for randomised controlled trial. *Trials*, 2020. 21: p. 1-3.
11. Dey, P., et al., Promoting uptake of influenza vaccination among health care workers: a randomized controlled trial. *Journal of public health medicine*, 2001: p. 346-348.
12. Looijmans-Van Den Akker, I., et al., Effects of a multi-faceted program to increase influenza vaccine uptake among health care workers in nursing homes: A cluster randomised controlled trial. *Vaccine*, 2010. 28(31): p. 5086-5092.
13. Upton, C.M., et al., Safety and efficacy of BCG re-vaccination in relation to COVID-19 morbidity in healthcare workers: A double-blind, randomised, controlled, phase 3 trial. *EClinicalMedicine*, 2022. 48.
14. Barbara, M., et al., The effect of giving influenza vaccination to general practitioners: a controlled trial [NCT002216761]. *BMC medicine*, 2006. 4(1): p. 1-10.
15. Bridges, C.B., et al., Effectiveness and cost-benefit of influenza vaccination of healthy working adults: a randomized controlled trial. *Jama*, 2000. 284(13): p. 1655-1663.
16. Pittet, L.F., et al., Randomized trial of BCG vaccine to protect against Covid-19 in health care workers. *New England Journal of Medicine*, 2023. 388(17): p. 1582-1596.
17. Borgey, F., et al., Effectiveness of an intervention campaign on influenza vaccination of professionals in nursing homes: A cluster-randomized controlled trial. *Vaccine*, 2019. 37(10): p. 1260-1265.
18. Imai, C., et al., A systematic review and meta-analysis of the direct epidemiological and economic effects of seasonal influenza vaccination on healthcare workers. *Plos one*, 2018. 13(6): p. e0198685.

19. Jacek JEdrzejek, M. and A. Mastalerz-Migas, Seasonal influenza vaccination of healthcare workers: A narrative review. *International Journal of Occupational Medicine and Environmental Health*, 2022. 35(2): p. 127.
20. Vorsters, A., et al., Overcoming barriers in HPV vaccination and screening programs. *Papillomavirus Research*, 2017. 4: p. 45-53.
21. Ng, A. and C. Lai, Effectiveness of seasonal influenza vaccination in healthcare workers: a systematic review. *Journal of Hospital Infection*, 2011. 79(4): p. 279-286.
22. Dini, G. , et al., Influenza vaccination in healthcare workers: A comprehensive critical appraisal of the literature. *Human vaccines & immunotherapeutics*, 2018. 14(3): p. 772-789.
23. Herzog, R. , et al., Are healthcare workers' intentions to vaccinate related to their knowledge, beliefs and attitudes? A systematic review. *BMC public health*, 2013. 13: p. 1-17.
24. Guillari, A., et al., Influenza vaccination and healthcare workers: barriers and predisposing factors. A literature review. *Acta Bio Medica: Atenei Parmensis*, 2021. 92(Suppl 2).
25. To, K., et al., Increasing the coverage of influenza vaccination in healthcare workers: review of challenges and solutions. *Journal of Hospital Infection*, 2016. 94(2): p. 133-142.

Table (1): Effectiveness of Healthcare Workers' Vaccination Programs: RCTs Findings Across Infectious Diseases

Study	Sample	Health professionals	Intervention	Effectiveness	Conclusions
RCI-	520	Emergency department personnel	Influenza vaccination	RR: 0.7 (0.6-0.9)	Significantly reduced influenza incidence. Vaccination Coverage reached 80%, highlighting feasibility. Subgroup analysis revealed higher efficacy In younger age groups.

4	852	Intensive care unit staff	Influenza vaccination	RR: 0.8 (0.7-1.0)	Moderate effectiveness in preventing influenza. Despite a vaccination coverage rate of 75% , a mild influenza outbreak occurred in the control group. Subsequent analysis revealed higher efficacy in vaccinated individuals with prior immunity.
---	-----	---------------------------	-----------------------	-------------------	---

<p>RCT-</p> <p>5</p>	<p>4,595</p>	<p>Primary care providers</p>	<p>Corona vaccination</p>	<p>RR: 0.4 (0.3-0.5)</p>	<p>Significant reduction in COVID-19 cases.</p> <p>Vaccination demonstrated a 60% reduction in symptomatic COVID-19 cases. Subgroup analysis by age and comorbidities revealed consistent efficacy.</p>
<p>Rer-</p> <p>6</p>	<p>657</p>	<p>Pediatric healthcare workers</p>	<p>TB vaccination</p>	<p>RR: 0.9 (0.8-1.1)</p>	<p>Limited effectiveness against TB infections. Despite high vaccination coverage, TB incidence reduction did not reach statistical significance. Subgroup analysis by age and years of service did not reveal substantial variations.</p>

7	3,376	Long-term care facility staff	Influenza vaccination	RR; 0.7 (0.6-0.8)	<p>Substantial reduction in influenza cases.</p> <p>Vaccination coverage of 85% correlated with a 40% decrease in influenza cases.</p> <p>Subgroup analysis by staff roles indicated higher efficacy in direct patient Care roles.</p>
8	1,812	Hospitalbased professionals	Corona vaccination	RR: 0.3 (0.2-0.4)	<p>Highly effective against COVID-19 infections.</p> <p>Vaccination demonstrated an 70% reduction in COVID-19 cases. Subgroup analysis by age and comorbidities consistentl showed hi h effica</p>

<p>RCT-</p> <p>9</p>	<p>2,534</p>	<p>Community health workers</p>	<p>TB vaccination</p>	<p>RR: 1.0 (0.9-1.1)</p>	<p>No significant impact on TB infection rates. Despite high vaccination coverage, TB incidence remained stable. Subgroup analysis by geographical location and years of service did not reveal substantial variations.</p>
----------------------	--------------	---------------------------------	-----------------------	--------------------------	---